

BELARUS NATIONAL ACADEMY OF SCIENCES
B.I. STEPANOV's INSTITUTE OF PHYSICS
BELARUSSIAN STATE UNIVERSITY

A. A. Bogush, V.M. Red'kov¹, G.G. Krylov²

SCHRÖDINGER PARTICLE IN MAGNETIC AND ELECTRIC FIELDS
IN LOBACHEVSKY AND RIEMANN SPACES

Schrödinger equation in Lobachevsky and Riemann 4-spaces has been solved in the presence of external magnetic field that is an analog of a uniform magnetic field in the flat space. Generalized Landau levels have been found, modified by the presence of the space curvature. In Lobachevsky 4-model the energy spectrum contains discrete and continuous parts, the number of bound states is finite; in Riemann 4-model all energy spectrum is discrete. Generalized Landau levels are determined by three parameters, the magnitude of the magnetic field B , the curvature radius ρ and the magnetic quantum number m . It has been shown that in presence of an additional external electric field the energy spectrum in the Riemann model can be also obtained analytically.

Pacs: 1130, 0230, 0365

keywords: magnetic field, Landau system, Schrödinger equation, hyperbolical 3-space, spherical 3-space, energy levels

Published:

Bogush A.A., Red'kov V.M., Krylov G.G.. Schrödinger particle in magnetic and electric fields in Lobachevsky and Riemann spaces. Nonlinear Phenomena in Complex Systems. 2008. Vol. 11. no 4, P. 403 – 416.

¹redkov@dragon.bas-net.by

²krylov@bsu.by

1 Introduction. Spaces of a constant curvature and uniform magnetic field

Quantum-mechanical Kepler problems in spaces of constant positive and negative curvature were formulated and solved many years ago by Schrödinger, Infeld, Schild, and Stefenson [1, 2, 3, 4]. Dynamical symmetry for quantum Kepler problems in spaces of constant curvature was investigated in [5, 6, 7, 8]. These models were used to describe bound state in nuclear physics [10, 11, 12] and nano-physics [13, 14].

In the present paper the non-relativistic quantum mechanical problems for a particle in a magnetic field on the background of hyperbolic Lobachevsky and spherical Riemann spaces are solved.

We start with Olevsky paper [15] where all 3-orthogonal coordinate systems (x^1, x^2, x^3) in 3-space of constant curvature, negative and positive, permitting the full separation of variables in the extended Helmholtz equation $(\Delta_2 + \lambda)f(x) = 0$ have been found. An idea consists in searching for an analog of cylindrical coordinate of Euclid space E_3 , in which one could easily define the concept of a uniform magnetic field in spaces H_3 and S_3 . Such uniform magnetic fields should be solutions of the corresponding Maxwell equations. Then we will try to examine the Landau problem [16, 18] of a Schrödinger's particle in the external magnetic field on the background of curved space models, H_3 and S_3 . Also, in present time we can note growing interest to Landau problem in the context of development in quantum mechanics in noncommutative space [24, 25, 26]. In the literature, the Landau problem was examined for Lobachevsky and spherical planes [19, 20, 21, 22, 23]; present paper agrees with the known results and adds to them.

In [15] in the list of coordinate systems, one can see the system number XI defined in space S_3 by the relations

$$\begin{aligned} dS^2 &= c^2 dt^2 - [\cos^2 z (dr^2 + \sin^2 r d\phi^2) - dz^2], \\ z &\in [-\pi/2, +\pi/2], \quad r \in [0, \pi], \quad \phi \in [0, 2\pi], \\ u_1 &= \cos z \sin r \cos \phi, \quad u_2 = \cos z \sin r \sin \phi, \\ u_3 &= \sin z, \quad u_0 = \cos z \cos r, \quad u_1^2 + u_2^2 + u_3^2 + u_0^2 = 1; \end{aligned}$$

and in space H_3 as

$$\begin{aligned} dS^2 &= c^2 dt^2 - [\cosh^2 z (dr^2 + \sinh^2 r d\phi^2) - dz^2], \\ z &\in (-\infty, +\infty), \quad r \in [0, +\infty], \quad \phi \in [0, 2\pi], \\ u_1 &= \cosh z \sinh r \cos \phi, \quad u_2 = \cosh z \sinh r \sin \phi, \\ u_3 &= \sinh z, \quad u_0 = \cosh z \cosh r, \quad u_1^2 + u_2^2 + u_3^2 - u_0^2 = -1. \end{aligned}$$

All coordinate u_a are dimensionless, what is achieved by dividing on the curvature radius ρ . When $\rho \rightarrow \infty$, these coordinates reduce to usual cylindrical ones in the space E_3

$$dS^2 = c^2 dt^2 - (dr^2 + r^2 d\phi^2 + dz^2).$$

Let us use the known form of the vector potential of the uniform constant magnetic field [2] in the flat space:

$$\mathbf{A} = \frac{1}{2} \mathbf{B} \times \mathbf{r}, \quad \mathbf{B} = (0, 0, B), \quad A^a = \frac{Br}{2} (0, -\sin \phi, \cos \phi, 0), \quad (1)$$

after translating to cylindrical coordinates

$$A_t = 0, \quad A_r = 0, \quad A_z = 0, \quad A_\phi = -\frac{Br^2}{2}. \quad (2)$$

Correspondingly we have a single non-vanishing component of the strength tensor which obeys the Maxwell equations:

$$F_{\phi r} = \partial_\phi A_r - \partial_r A_\phi = Br , \quad \frac{1}{r} \frac{\partial}{\partial r} r F^{\phi r} = \frac{1}{r} \frac{\partial}{\partial r} r \left(-\frac{1}{r^2} \right) Br \equiv 0 . \quad (3)$$

Such description of a uniform magnetic field in cylindrical coordinates may be extended quite easily to curved space models. Let us start with the following potential in the spherical space S_3

$$A_\phi = 2B (\cos r - 1) , \quad F_{\phi r} = B \sin r , \quad (4)$$

Analogously, in Lobachevsky model we start with H_3

$$A_\phi = -2B (\cosh r - 1) , \quad F_{\phi r} = B \sinh r . \quad (5)$$

As one can see both potentials are solutions of the respective Maxwell equations, and in the vanishing curvature limit they are reduced to (3).

2 Separation of variables in the Schrödinger equation

Non-relativistic Schrödinger equation in a space-time with the metrics

$$dS^2 = (dx^0)^2 + g_{kl}(x) dx^k dx^l , \quad g = \det (g_{kl}(x))$$

has the form

$$(i\hbar \partial_t + e A_0) \Psi = H \Psi ,$$

$$H = \frac{-1}{2M} \left[\left(\frac{i\hbar}{\sqrt{-g}} \partial_k \sqrt{-g} + \frac{e}{c} A_k \right) g^{kl} \left(i\hbar \partial_l + \frac{e}{c} A_l \right) \right] \Psi .$$

In the space S_3 , parameterized by cylindric coordinates and in the presence of the external uniform magnetic field, the Hamiltonian H takes the form

$$H = \frac{\hbar^2}{2MR^2} \left[-\frac{1}{\cos^2 z \sin r} \frac{\partial}{\partial r} \sin r \frac{\partial}{\partial r} + \frac{1}{\cos^2 z \sin^2 r} \left(i \frac{\partial}{\partial \phi} + \frac{e}{\hbar c} A_\phi \right)^2 - \frac{1}{\cos^2 z} \frac{\partial}{\partial z} \cos^2 z \frac{\partial}{\partial z} \right]$$

and the variables may be separated by the substitution:

$$\Psi = e^{-iEt/\hbar} e^{im\phi} Z(z) R(r) , \quad \epsilon = E/(\hbar^2/M\rho^2) , \quad (6)$$

$$\frac{1}{Z} \left(2\epsilon \cos^2 z Z + \frac{d}{dz} \cos^2 z \frac{dZ}{dz} \right)$$

$$= \frac{1}{R} \left(-\frac{1}{\sin r} \frac{d}{dr} \sin r \frac{dR}{dr} + \frac{1}{\sin^2 r} \left(-m + \frac{e}{\hbar c} A_\phi \right)^2 R \right) .$$

Introducing the separation constant λ , we arrive at

$$\frac{d}{dz} \cos^2 z \frac{dZ}{dz} + 2\epsilon \cos^2 z Z = \lambda Z , \quad (7)$$

$$-\frac{1}{\sin r} \frac{d}{dr} \sin r \frac{dR}{dr} + \frac{1}{\sin^2 r} \left(m + \frac{eB}{\hbar c} 2\rho^2 \sin^2 \frac{r}{2} \right)^2 R = \lambda R . \quad (8)$$

In the same manner, in the Lobachevsky space H_3 we arrive at

$$\frac{d}{dz} \cosh^2 z \frac{dZ}{dz} + 2\epsilon \cosh^2 z Z = \lambda Z, \quad (9)$$

$$-\frac{1}{\sinh r} \frac{d}{dr} \sinh r \frac{dR}{dr} + \frac{1}{\sinh^2 r} \left(m + \frac{eB}{\hbar c} 2\rho^2 \sinh^2 \frac{r}{2} \right)^2 R = \lambda R. \quad (10)$$

In the limit of flat space these equations become more simple:

$$\begin{aligned} \frac{d^2 Z}{dz^2} + \frac{2EM}{\hbar^2} Z &= \Lambda Z, \quad Z = e^{\pm i P z / \hbar}, \quad \Lambda = \frac{2M}{\hbar^2} \left(E - \frac{P^2}{2M} \right), \\ -\frac{1}{r} \frac{d}{dr} r \frac{dR}{dr} + \frac{1}{r^2} \left(m + \frac{eB}{\hbar c} \frac{r^2}{2} \right)^2 R &= \frac{2M}{\hbar^2} \left(E - \frac{P^2}{2M} \right) R \end{aligned} \quad (11)$$

which coincide with the known one [18]:

$$\frac{\hbar^2}{2M} \left(\frac{d^2 R}{dr^2} - \frac{1}{r} \frac{dR}{dr} - \frac{m^2}{r^2} R \right) + \left[E - \frac{P^2}{2M} - \frac{\omega^2}{8} r^2 - \frac{\hbar \omega m}{2} \right] R = 0$$

with $\omega = eB/Mc$.

3 Radial solutions in space H_3

Below the notion $\frac{eB}{\hbar c} \rho^2 \implies B$ will be used, then eq. (10) reads

$$\frac{d^2}{dr^2} R + \frac{1}{\tanh r} \frac{dR}{dr} - \frac{1}{\sinh^2 r} [m + B(\cosh r - 1)]^2 R + \lambda R = 0. \quad (12)$$

In new variables

$$\cosh r - 1 = -2z, \quad z = -\sinh^2 \frac{r}{2} \in (-\infty, 0],$$

for the radial equation we get

$$\left[z(1-z) \frac{d^2}{dz^2} + (1-2z) \frac{d}{dz} - \frac{1}{4} \left(\frac{m^2}{z} - 4B^2 + \frac{(m-2B)^2}{1-z} \right) - \lambda \right] R = 0. \quad (13)$$

Let us specify the behavior of the function R near to the points 0 and 1 (take note that the value 1 does not belong to the physical domain of the coordinate r):

$$\begin{aligned} \text{if } z \longrightarrow 0, \text{ then } R(z) &\sim z^a, \quad a = \pm \frac{m}{2}; \\ \text{if } z \longrightarrow 1, \text{ then } R(z) &\sim (1-z)^b, \quad b = \pm \frac{m-2B}{2}. \end{aligned}$$

Making substitution $R = z^a (1-z)^b F$, from eq. (13) we get

$$\begin{aligned} &+ z(1-z) F'' + [a(1-z) - bz + a(1-z) - bz + (1-2z)] F' + \\ &+ \frac{1}{z} [a(a-1) + a - \frac{m^2}{4}] F + \frac{1}{1-z} [b(b-1) + b - \frac{(m-2B)^2}{4}] F - \\ &- [a(a+1) + 2ab + b(b+1) - B^2 + \lambda] F = 0; \end{aligned} \quad (14)$$

Imposing conditions $a = \pm m/2$, $b = \pm(m - 2B)/2$ we obtain

$$\begin{aligned} R &= z^a(1-z)^b F, \\ z(1-z)F'' + [(2a+1) - 2(a+b+1)z]F' \\ &- [a(a+1) + 2ab + b(b+1) - B^2 + \lambda]F = 0 \end{aligned}$$

which is the equation of hypergeometric type: $z(1-z)F + [\gamma - (\alpha + \beta + 1)z]F' - \alpha\beta F = 0$. Thus, the radial function is constructed in hypergeometric functions as follows: (to obtain the bound states we must take the parameters a and b positive and negative respectively)

$$\begin{aligned} z &= -\sinh^2 \frac{r}{2}, z \in (-\infty, +0], \\ R &= (-\sinh \frac{r}{2})^{|m|} (\cosh \frac{r}{2})^{-|m-2B|} {}_2F_1(\alpha, \beta; \gamma; -\sinh^2 \frac{r}{2}) \end{aligned} \quad (15)$$

where (α, β) are defined by

$$\alpha + \beta = 2a + 2b + 1, \quad \alpha\beta = (a+b)(a+b+1) - B^2 + \lambda$$

that is

$$\begin{aligned} a &= +\frac{|m|}{2}, \quad b = -\frac{|m-2B|}{2}, \\ \alpha &= a + b + \frac{1}{2} - \sqrt{B^2 + \frac{1}{4} - \lambda}, \\ \beta &= a + b + \frac{1}{2} + \sqrt{B^2 + \frac{1}{4} - \lambda}, \\ \gamma &= 2a + 1 = +|m| + 1. \end{aligned} \quad (16)$$

The first possibility to obtain polynomial is

$$\alpha = a + b + \frac{1}{2} - \sqrt{B^2 + \frac{1}{4} - \lambda} = -n = 0, -1, \dots$$

from whence it follows the quantization rule

$$a + b + \frac{1}{2} + n \geq 0, \quad \lambda = \frac{1}{4} + B^2 - \left(a + b + n + \frac{1}{2}\right)^2 \quad (17)$$

or

$$\begin{aligned} \frac{|m|}{2} - \frac{|m-2B|}{2} + n + \frac{1}{2} &\geq 0, \\ \lambda &= \frac{1}{4} + B^2 - \left(\frac{|m|}{2} - \frac{|m-2B|}{2} + n + \frac{1}{2}\right)^2, \end{aligned} \quad (18)$$

$$R = (-\sinh \frac{r}{2})^{|m|} (\cosh \frac{r}{2})^{-|m-2B|} {}_2F_1(-n, |m| - |m-2B| + 1 + n; |m| + 1, -\sinh^2 \frac{r}{2}). \quad (19)$$

The second possibility to obtain the polynomial is

$$\beta = a + b + \frac{1}{2} + \sqrt{B^2 + \frac{1}{4} - \lambda} = -n$$

that is

$$a + b + \frac{1}{2} + n \leq 0, \quad \lambda = \frac{1}{4} + B^2 - (a + b + \frac{1}{2} + n)^2 \quad (20)$$

or

$$\begin{aligned} \frac{|m|}{2} - \frac{|m-2B|}{2} + n + \frac{1}{2} &\leq 0, \\ \lambda &= \frac{1}{4} + B^2 - \left(\frac{|m|}{2} - \frac{|m-2B|}{2} + n + \frac{1}{2} \right)^2, \end{aligned} \quad (21)$$

$$R = (-\sinh \frac{r}{2})^{|m|} (\cosh \frac{r}{2})^{-|m-2B|} {}_2F_1(|m| - |m-2B| + 1 + n, -n; |m| + 1; -\sinh^2 \frac{r}{2}). \quad (22)$$

Now, let we assume that $B > 0$. There exists five possibilities:

$$m = 0, m < 0, 0 < m < 2B, m > 2B, m = 2B.$$

For the case $m = 0$ we have $a = 0$, $b = -B$. Applying the quantization condition (18) we get a very special situation: namely, one separate bound state may exist or not depending on the value of a magnetic field:

$$\begin{aligned} 0 < B - n &\leq \frac{1}{2}, \quad \lambda - \frac{1}{4} = 2B(n + 1/2) - (n + 1/2)^2, \\ R &= (\cosh \frac{r}{2})^{-2B} {}_2F_1(-n, -2B + 1 + n; +1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim (\cosh \frac{r}{2})^{-2B} (\sinh^2 \frac{r}{2})^n \sim e^{-(B-n)r}. \end{aligned} \quad (23)$$

Applying the quantization rule (21) we get a finite series of bound states:

$$\begin{aligned} n + \frac{1}{2} &\leq B, \quad n = 0, 1, 2, \dots, N_B, \quad \lambda - 1/4 = 2B(n + 1/2) - (N + 1/2)^2, \\ R &= (\cosh \frac{r}{2})^{-2B} {}_2F_1(-2B + 1 + n, -n; |m| + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim (\cosh \frac{r}{2})^{-2B} (\sinh^2 \frac{r}{2})^n \sim e^{-(B-n)r}. \end{aligned} \quad (24)$$

Now let us consider the case

$$m < 0, \quad a = -\frac{m}{2} > 0, \quad b = \frac{m}{2} - B < 0.$$

Applying the quantization condition (18) we again get a special situation: one separate bound state may exist or not depending on the value of magnetic field:

$$\begin{aligned} 0 < B - n &\leq \frac{1}{2}, \quad \lambda - \frac{1}{4} = 2B(n + 1/2) - (n + 1/2)^2, \\ R &= (-\sinh \frac{r}{2})^{-m} (\cosh \frac{r}{2})^{m-2B} {}_2F_1(-n, -2B + 1 + n; -m + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim e^{-(B-n)r}. \end{aligned} \quad (25)$$

Applying the quantization rule (21) we get a finite series of bound states:

$$\begin{aligned} n + \frac{1}{2} &\leq B, \quad n = 0, 1, 2, \dots, N_B, \\ \lambda - 1/4 &= 2B(n + 1/2) - (n + 1/2)^2, \\ R &= (-\sinh \frac{r}{2})^{-m} (\cosh \frac{r}{2})^{m-2B} {}_2F_1(-2B + 1 + n, -n; -m + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim e^{-(B-n)r}. \end{aligned} \quad (26)$$

Now let us consider the case

$$0 < m < 2B, \quad a = \frac{m}{2} > 0, \quad b = \frac{m}{2} - B < 0.$$

Applying the quantization condition (18) we get a very special situation: one separate bound state may exist or not depending on the value of magnetic field and magnetic quantum number:

$$\begin{aligned} 0 \leq B - m - n \leq \frac{1}{2}, \quad \lambda - 1/4 &= 2B(m + n + 1/2) - (m + n + 1/2)^2, \\ R &= (-\sinh \frac{r}{2})^m (\cosh \frac{r}{2})^{m-2B} {}_2F_1(-n, 2m - 2B + 1 + n; m + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim e^{-(B-m-n)r}. \end{aligned} \quad (27)$$

Applying the quantization rule (21) we get a finite series of bound states:

$$\begin{aligned} m + \frac{1}{2} + n &\leq B, \quad n = 0, 1, \dots, N_{B,m}, \\ \lambda - 1/4 &= 2B(m + n + 1/2) - (m + n + 1/2)^2, \\ R &= (-\sinh \frac{r}{2})^m (\cosh \frac{r}{2})^{m-2B} {}_2F_1(2m - 2B + 1 + n, -n; m + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim e^{-(B-m-n)r}. \end{aligned} \quad (28)$$

For the case

$$m > 2B \quad a = \frac{m}{2}, \quad b = -\frac{m - 2B}{2} < 0,$$

with the rule (18) we get

$$\begin{aligned} B + \frac{1}{2} + n &\geq 0, \quad \lambda - \frac{1}{4} = -2B(n + \frac{1}{2}) - (n + \frac{1}{2})^2, \\ R &= (-\sinh \frac{r}{2})^m (\cosh \frac{r}{2})^{-m+2B} {}_2F_1(-n, 2B + 1 + n; m + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim e^{(B+n)r} \rightarrow \infty; \end{aligned} \quad (29)$$

this is not the case of bound states. With the rule (21) we arrive at

$$B + \frac{1}{2} + n \leq 0 \quad (30)$$

that can not be fulfilled together with the quantization conditions whereas $n \geq 0$.

And finally for the last variant:

$$m = 2B > 0, \quad a = m = 2B, \quad b = 0,$$

we get

$$\begin{aligned} B + n + \frac{1}{2} &\geq 0, \quad \lambda - \frac{1}{4} = 2B(n + \frac{1}{2}) - (n + \frac{1}{2})^2, \\ R &= (-\sinh \frac{r}{2})^{2B} {}_2F_1(-n, |m| - |m - 2B| + 1 + n; |m| + 1; -\sinh^2 \frac{r}{2}), \\ R_{r \rightarrow +\infty} &\sim e^{(B+n)r} \rightarrow \infty \end{aligned}$$

which is not a bound state. Quantization according to (21) cannot provide us with bound states because of relationship $B + n + 1/2 > 0$.

Let us collect the results together:

separate states: that may be presented by the single formula:

$$m < 2B, \quad n < B \leq n + 1/2, \quad n = 0, 1, 2, \dots, \\ \lambda - \frac{1}{4} = 2B \left(\frac{m + |m|}{2} + n + 1/2 \right) - \left(\frac{m + |m|}{2} + n + \frac{1}{2} \right)^2; \quad (31)$$

series of states: given by a formula

$$m < 2B, \quad \frac{m + |m|}{2} + n + 1/2 \leq B, \\ \lambda - 1/4 = 2B \left(\frac{m + |m|}{2} + n + 1/2 \right) - \left(\frac{m + |m|}{2} + n + \frac{1}{2} \right)^2, \quad n = 0, 1, \dots, N_B. \quad (32)$$

In usual unites, the last relation reads

$$\lambda - \frac{1}{4} = \rho^2 \lambda_0 - \frac{1}{4}, \quad \lim_{\rho \rightarrow \infty} \lambda_0 = \frac{2M}{\hbar^2} \left(E - \frac{P^2}{2M} \right), \quad m < 2B, \quad m + n + 1/2 \leq \frac{eB}{\hbar c} \rho^2, \\ \rho^2 \lambda_0 - \frac{1}{4} = 2 \frac{eB}{\hbar c} \rho^2 \left(\frac{m + |m|}{2} + n + 1/2 \right) - \left(\frac{m + |m|}{2} + n + 1/2 \right)^2, \quad n = 0, 1, \dots, N_B. \quad (33)$$

In the limit of a flat space, from (33) the known result follows:

$$E - \frac{P^2}{2M} = \frac{eB\hbar}{Mc} \left(\frac{m + |m|}{2} + n + 1/2 \right).$$

For the case of the positive orientation of magnetic field $B < 0$, we should take into account the symmetry provided by initial differential equation: $m \rightarrow m' = -m$, $B \rightarrow B' = -B$.

4 Differential equation for $Z(z)$ in H_3

Now, let us examine the differential equation for $Z(z)$ in space H_3 :

$$\frac{d^2 Z}{dz^2} + 2 \frac{\sinh z}{\cosh z} \frac{dZ}{dz} + 2\epsilon Z - \frac{\lambda}{\cosh^2 z} Z = 0. \quad (34)$$

Changing the variable as $\sinh z = i(2x - 1)$, we get

$$x(1-x) \frac{d^2 Z}{dx^2} - 3(2x-1) \frac{1}{2} \frac{dZ}{dx} - \left(2\epsilon - \frac{\lambda}{4x(1-x)} \right) Z = 0.$$

Taking the function in the form $Z = x^a(1-x)^b F$, we obtain

$$+ x(1-x) F'' + [2a(1-x) - 2bx - 3x + 3/2] F' + \left[a(a-1) + \frac{3a}{2} + \frac{\lambda}{4} \right] \frac{1}{x} F \\ + \left[b(b-1) + 3b - \frac{3b}{2} + \frac{\lambda}{4} \right] \frac{1}{1-x} F + [-a(a-1) - ab - ab - b(b-1) - 3a - 3b - 2\epsilon] F = 0. \quad (35)$$

Imposing additional restrictions

$$a^2 + \frac{a}{2} + \frac{\lambda}{4} = 0, \quad a = \frac{-1 \pm \sqrt{1-4\lambda}}{4}, \quad b^2 + \frac{b}{2} + \frac{\lambda}{4} = 0, \quad b = \frac{-1 \pm \sqrt{1-4\lambda}}{4}, \quad (36)$$

we arrive at more simple equation

$$x(1-x)F'' + [(2a+3/2) - x(2a+2b+3)]F' - [(a+b+1)^2 + 2\epsilon - 1]F = 0 \quad (37)$$

which is of hypergeometric type

$$x(1-x)F + [C - (A+B+1)x]F' - ABF = 0$$

with parameters defined by

$$A+B = 2(a+b+1), \quad AB = (a+b+1)^2 + 2\epsilon - 1.$$

That is

$$\begin{aligned} A &= a+b+1 + \sqrt{1-2\epsilon}, & B &= a+b+1 - \sqrt{1-2\epsilon}, & C &= 2a+3/2; \\ Z &= x^a(1-x)^b {}_2F_1(A, B; C; x), & x &= \frac{1 - \imath \sinh z}{2}, & (1-x) &= \frac{1 + \imath \sinh z}{2}, \\ a &= \frac{-1 \pm \sqrt{1-4\lambda}}{4}, & b &= \frac{-1 \pm \sqrt{1-4\lambda}}{4}. \end{aligned} \quad (38)$$

To find behavior of these solutions at infinity, one can use the Kummer's relations [27]

$$\begin{aligned} U_1 &= {}_2F_1(A, B; C; x), \\ U_3 &= (-x)^{-A} {}_2F_1(A, A-C+1; A-B+1; x^{-1}), \\ U_4 &= (-x)^{-B} {}_2F_1(B-C+1, B; B-A+1; x^{-1}), \\ U_1 &= \frac{\Gamma(C)\Gamma(B-A)}{\Gamma(C-A)\Gamma(B)} U_3 + \frac{\Gamma(C)\Gamma(A-B)}{\Gamma(C-B)\Gamma(A)} U_4. \end{aligned} \quad (39)$$

Then

$$\begin{aligned} Z &= x^a(1-x)^b U_1 = \\ &= \frac{\Gamma(C)\Gamma(B-A)}{\Gamma(C-A)\Gamma(B)} x^a(1-x)^b (-x)^{-(a+b+1+i\sqrt{2\epsilon-1})} {}_2F_1(A, A-C+1; A-B+1; x^{-1}) \\ &+ \frac{\Gamma(C)\Gamma(A-B)}{\Gamma(C-B)\Gamma(A)} x^a(1-x)^b (-x)^{-(a+b+1-i\sqrt{2\epsilon-1})} {}_2F_1(B-C+1, B; B-A+1; x^{-1}). \end{aligned} \quad (40)$$

Therefore, asymptotically all three solutions U_1, U_3, U_4 vanish when $x \rightarrow \infty$:

$$\begin{aligned} Z &= x^a(1-x)^b U_1 \sim \\ &\sim \frac{\Gamma(C)\Gamma(B-A)}{\Gamma(C-A)\Gamma(B)} x^a(1-x)^b (-x)^{-a-b-1-i\sqrt{2\epsilon-1}} + \frac{\Gamma(C)\Gamma(A-B)}{\Gamma(C-B)\Gamma(A)} x^a(1-x)^b \\ &\quad \times (-x)^{-a-b-1+i\sqrt{2\epsilon-1}} = \\ &(-1)^a \left[\frac{\Gamma(C)\Gamma(B-A)}{\Gamma(C-A)\Gamma(B)} (-x)^{-1-i\sqrt{2\epsilon-1}} + \frac{\Gamma(C)\Gamma(A-B)}{\Gamma(C-B)\Gamma(A)} (-x)^{-1+i\sqrt{2\epsilon-1}} \right] \rightarrow 0. \end{aligned} \quad (41)$$

Let $a = b$, then

$$\begin{aligned}
a = b &= \frac{-1 \pm 2i\sqrt{\lambda - 1/4}}{4}, \quad a + b = \frac{-1 \pm 2i\sqrt{\lambda - 1/4}}{2}, \\
Z &= x^a(1-x)^a {}_2F_1(A, B; C; x) = \left(\frac{\cosh^2 z}{4}\right)^a {}_2F_1(A, B; C; x), \\
A &= \frac{1}{2} + i(\sqrt{2\epsilon - 1} \pm \sqrt{\lambda - \frac{1}{4}}), \quad B = \frac{1}{2} - i(\sqrt{2\epsilon - 1} \mp \sqrt{\lambda - \frac{1}{4}}), \\
C &= 1 \pm i\sqrt{\lambda - 1/4}.
\end{aligned} \tag{42}$$

One may note that from (42) it follows

$$(A - \frac{1}{2})(B - \frac{1}{2}) = (2\epsilon - 1) - (\lambda - 1/4).$$

It has sense to introduce a new parameter

$$(2\epsilon - 1) - (\lambda - 1/4) = \nu^2, \quad \text{or} \quad (\lambda - 1/4) = (2\epsilon - 1) - \nu^2 \tag{43}$$

which generalizes the known relation in the flat space model $\lambda = 2\epsilon - k^2$. It is the matter of simple calculation to determine the operator with eigenvalues $\nu^2 + 3/4$:

$$\hat{\nu}^2 + \frac{3}{4} = 2\hat{H}(r, \phi, z) - \hat{\lambda}(r).$$

Indeed, with the notion

$$\begin{aligned}
\hat{\lambda}(r, \phi) &= -\frac{1}{\sinh r} \frac{\partial}{\partial r} \sinh r \frac{\partial}{\partial r} + \frac{1}{\sinh^2 r} \left[i \frac{\partial}{\partial \phi} - B(\cosh r - 1) \right]^2, \\
2\hat{H}(r, \phi, z) &= \frac{1}{\cosh^2 z} \hat{\lambda}(r, \phi) - \frac{1}{\cosh^2 z} \frac{\partial}{\partial z} \cosh^2 z \frac{\partial}{\partial z}.
\end{aligned} \tag{44}$$

We get

$$\hat{\nu}^2(r, \phi, z) + \frac{3}{4} = \left(\frac{1}{\cosh^2 z} - 1 \right) \hat{\lambda}(r, \phi) - \frac{1}{\cosh^2 z} \frac{\partial}{\partial z} \cosh^2 z \frac{\partial}{\partial z}. \tag{45}$$

and the commutative relations $[2H, \hat{\lambda}(r)] = 0$ holds.

5 Radial equation in spherical space

Now let us consider the radial equation arising in spherical space. With the use of dimensionless parameter $\frac{eB}{\hbar c} \rho^2 \implies B$ the equation reads

$$\frac{d^2}{dr^2} R + \frac{1}{\tan r} \frac{dR}{dr} - \frac{1}{\sin^2 r} [m + B(1 - \cos r)]^2 R + \lambda R = 0. \tag{46}$$

With variable change given by $1 - \cos r = 2z$ we have

$$[z(1-z)\frac{d^2}{dz^2} + (1-2z)\frac{d}{dz} - \frac{1}{4}(\frac{m^2}{z} - 4B^2 + \frac{(m+2B)^2}{1-z}) + \lambda]R = 0. \quad (47)$$

After substitution $R = z^a(1-z)^bF$ we produce

$$\begin{aligned} & z(1-z)F'' + [a(1-z) - bz + a(1-z) - bz + (1-2z)]F' + \frac{1}{z}[a(a-1) + a - m^2/4]F \\ & + \frac{1}{1-z}[b(b-1) + b - (m+2B)^2/4]F - [a(a+1) + 2ab + b(b+1) - B^2 - \lambda]F = 0. \end{aligned} \quad (48)$$

Imposing the conditions $a = \pm m/2$, $b = \pm(m+2B)/2$ we arrive at

$$z(1-z)F'' + [(2a+1) - 2(a+b+1)z]F' - [a(a+1) + 2ab + b(b+1) - B^2 - \lambda]F = 0 \quad (49)$$

which is of hypergeometric type. Thus, the problem is solved as

$$\begin{aligned} & z = \sin^2 \frac{r}{2}, \quad z \in [0, +1], \quad r \in [0, +\pi]; \\ & R = (\sin \frac{r}{2})^{+|m|} (\cos \frac{r}{2})^{+|m+2B|} {}_2F_1(\alpha, \beta; \gamma; -\sin^2 \frac{r}{2}) \end{aligned} \quad (50)$$

where

$$\begin{aligned} & a = +\frac{|m|}{2}, \quad b = +\frac{|m+2B|}{2}, \quad \alpha = a + b + \frac{1}{2} - \sqrt{B^2 + \frac{1}{4} + \lambda}, \\ & \beta = a + b + \frac{1}{2} + \sqrt{B^2 + \frac{1}{4} + \lambda}, \quad \gamma = +|m| + 1. \end{aligned} \quad (51)$$

The quantization condition is given by

$$\alpha = a + b + \frac{1}{2} - \sqrt{B^2 + \frac{1}{4} + \lambda} = -n, \quad n = 0, -1, \dots$$

from whence it follows relations for possible bound states:

$$\begin{aligned} & \lambda + \frac{1}{4} = -B^2 + (a + b + \frac{1}{2} + n)^2, \quad n = 0, -1, -2, \dots \\ & R = (\sin \frac{r}{2})^{+|m|} (\cos \frac{r}{2})^{+|m+2B|} {}_2F_1(-n, |m| + |m+2B| + 1 + n; |m| + 1; -\sin^2 \frac{r}{2}), \end{aligned} \quad (52)$$

While examining wave functions in spherical space one must take into account peculiarities in parameterization of the space S_3 by the coordinates (r, ϕ, z) :

$$\begin{aligned} & z \in [-\pi/2, +\pi/2], \quad r \in [0, \pi], \quad \phi \in [0, 2\pi], \\ & u_1 = \cos z \sin r \cos \phi, \quad u_2 = \cos z \sin r \sin \phi, \\ & u_3 = \sin z, \quad u_0 = \cos z \cos r, \quad u_1^2 + u_2^2 + u_3^2 + u_0^2 = 1. \end{aligned} \quad (53)$$

In particular, we have

$$\begin{aligned} & r = 0, \quad u_1 = 0, \quad u_2 = 0, \quad u_3 = \sin z, \quad u_0 = +\cos z, \\ & r = \pi, \quad u_1 = 0, \quad u_2 = 0, \quad u_3 = \sin z, \quad u_0 = -\cos z, \\ & z \in [-\pi/2, +\pi/2] \end{aligned} \quad (54)$$

which means that the full closed curve $u_0^2 + u_3^2 = 1$ in S_3 is parameterized by two pieces:

$$\{(r = 0, \phi - \text{mute}, z) + (r = \pi, \phi - \text{mute}, z)\}.$$

Correspondingly, when $m \neq 0$, the function $R(r)$ must vanish in $r = 0, \pi$:

$$m \neq 0, \quad R(0) = 0, \quad R(\pi) = 0. \quad (55)$$

First, let us examine the case $m = 0$, then

$$\begin{aligned} a &= 0, \quad b = +B, \\ \lambda + \frac{1}{4} &= 2B(n + 1/2) + (n + 1/2)^2, \\ R &= (\cos \frac{r}{2})^{+2B} {}_2F_1(-n, 2B + n + 1; 1; -\sin^2 \frac{r}{2}), \\ R_{r \rightarrow 0} &= 1, \quad R_{r \rightarrow \pi} = 0. \end{aligned} \quad (56)$$

Because $m = 0$ and the wave function does not depend on ϕ , this solution is acceptable by continuity reasons. Now, let us examine the case $m > 0$, then

$$\begin{aligned} a &= +\frac{m}{2}, \quad b = +\frac{m + 2B}{2}, \\ \lambda + \frac{1}{4} &= 2B(n + m + 1/2) + (n + m + 1/2)^2, \\ R &= (\sin \frac{r}{2})^{+m} (\cos \frac{r}{2})^{m+2B} \\ &\times {}_2F_1(-n, 2B + 2m + n + 1; m + 1; -\sin^2 \frac{r}{2}), \\ R_{r \rightarrow 0} &= 0, \quad R_{r \rightarrow \pi} = 0; \end{aligned} \quad (57)$$

these are the correct single-valued solutions. Let us consider the variant $m < -2B$, then

$$\begin{aligned} a &= -\frac{m}{2}, \quad b = -\frac{m + 2B}{2} > 0, \\ \lambda + \frac{1}{4} &= -2B(n - m + 1/2) + (n - m + 1/2)^2 \\ &= (n - m + 1/2)((n - m + 1/2) - 2B) > 0, \\ R &= (\sin \frac{r}{2})^{-m} (\cos \frac{r}{2})^{-(m+2B)} \\ &\times {}_2F_1(-n, -2m - 2B + 1 + n; -m + 1; -\sin^2 \frac{r}{2}), \\ R_{r \rightarrow 0} &= 0, \quad R_{r \rightarrow \pi} = 0. \end{aligned} \quad (58)$$

Let us examine the case $-2B < m < 0$, then

$$\begin{aligned} a &= -\frac{m}{2} > 0, \quad b = \frac{m + 2B}{2} > 0, \\ \lambda + \frac{1}{4} &= 2B(n + 1/2) + (n + 1/2)^2, \\ R &= (\sin \frac{r}{2})^{-m} (\cos \frac{r}{2})^{m+2B} \\ &\times {}_2F_1(-n, 2B + n + 1; -m + 1; -\sin^2 \frac{r}{2}), \\ R_{r \rightarrow 0} &= 0, \quad R_{r \rightarrow \pi} = 0. \end{aligned} \quad (59)$$

And finally, the last variant is $m = -2B$, then

$$\begin{aligned}
a &= +B, \quad b = 0, \\
\lambda + \frac{1}{4} &= 2B(n + 1/2) + (n + 1/2)^2, \\
R &= (-\sin \frac{r}{2})^{+2B} \\
&\times {}_2F_1(-n, 2B + n + 1; -B + 1; -\sin^2 \frac{r}{2}), \\
R_{r \rightarrow +0} &= 0, \quad R_{r \rightarrow +\pi} = 1.
\end{aligned} \tag{60}$$

We faced here with the case of discontinues of wave functions in S_3 , because the wave function preserves dependence on ϕ at $r \rightarrow \pi$.

Collecting results together, we have

$$\begin{aligned}
m > 0, \quad \lambda + \frac{1}{4} &= (n + 1/2 + m)(n + 1/2 + m + 2B); \\
m < -2B, \quad \lambda + \frac{1}{4} &= (n + 1/2 - m)(n + 1/2 - m - 2B); \\
-2B < m \leq 0, \quad]\lambda + \frac{1}{4} &= (n + 1/2)(n + 1/2 - 2B).
\end{aligned} \tag{61}$$

In usual unites, these formulas look

$$\begin{aligned}
m > 0 \quad \rho^2 \lambda_0 + \frac{1}{4} &= +2 \frac{eB}{\hbar c} \rho^2 (n + m + 1/2) + (n + m + 1/2)^2; \\
m < -2 \frac{eB}{\hbar c} \rho^2 \quad \rho^2 \lambda_0 + \frac{1}{4} &= -2 \frac{eB}{\hbar c} \rho^2 (n - m + 1/2) + (n - m + 1/2)^2; \\
-2 \frac{eB}{\hbar c} \rho^2 < m \leq 0, \quad \rho^2 \lambda_0 + \frac{1}{4} &= 2 \frac{eB}{\hbar c} \rho^2 (n + 1/2) + (n + 1/2)^2.
\end{aligned} \tag{62}$$

When $\rho \rightarrow 0$, the case $m < -\infty$ turns to be vanishing, and two other cases give

$$\begin{aligned}
m < 0, \quad \lambda_0 &= 2 \frac{eB}{\hbar c} (n + 1/2), \\
m \geq 0, \quad \lambda_0 &= +2 \frac{eB}{\hbar c} (n + m + 1/2).
\end{aligned} \tag{63}$$

Thus, we arrive at the known result in the flat space model:

$$\lim_{\rho \rightarrow \infty} \lambda_0 = \frac{2M}{\hbar^2} \left(E - \frac{P^2}{2M} \right), \quad E - \frac{P^2}{2M} = \frac{eB\hbar}{Mc} \left(\frac{m + |m|}{2} + n + 1/2 \right). \tag{64}$$

6 Solutions of the equation for $Z(z)$ in the space H_3

Now, let us consider the differential equation for $Z(z)$ in space S_3 :

$$\frac{d^2 Z}{dz^2} - 2 \frac{\sin z}{\cos z} \frac{dZ}{dz} + 2\epsilon Z - \frac{\lambda}{\cos^2 z} Z = 0. \quad (65)$$

Changing the variable $\sin z = y$, we get

$$(1 - y^2) \frac{d^2 Z}{dy^2} - 3y \frac{dZ}{dy} + \left(2\epsilon - \frac{\lambda}{1 - y^2} \right) Z = 0. \quad (66)$$

In the variable $y = (2x - 1)$ it takes the form

$$x(1 - x) \frac{d^2 Z}{dx^2} - \frac{3}{2}(2x - 1) \frac{dZ}{dx} + \left(2\epsilon - \frac{\lambda}{4x(1 - x)} \right) Z = 0. \quad (67)$$

With the use of the substitution $Z = x^a(1 - x)^b F$, eq. (67) leads to

$$\begin{aligned} & + x(1 - x)F'' + [2a(1 - x) - 2bx - 3x + \frac{3}{2}]F' + [a(a - 1) + 3a/2 - \lambda/4]\frac{1}{x}F \\ & + [b(b - 1) + 3b - 3b/2 - \lambda/4]\frac{1}{1 - x}F + [-a(a - 1) - ab - ab - b(b - 1) - 3a - 3b + 2\epsilon]F = 0. \end{aligned} \quad (68)$$

Imposing the following restrictions

$$\begin{aligned} a^2 + \frac{a}{2} - \frac{\lambda}{4} &= 0, & a &= \frac{-1 \pm \sqrt{1 + 4\lambda}}{4}, \\ b^2 + \frac{b}{2} - \frac{\lambda}{4} &= 0, & b &= \frac{-1 \pm \sqrt{1 + 4\lambda}}{4} \end{aligned} \quad (69)$$

we reduce eq. (68) to

$$x(1 - x)F'' + [(2a + 3/2) - x(2a + 2b + 3)]F' - [(a + b + 1)^2 - 2\epsilon - 1]F = 0, \quad (70)$$

which is of hypergeometric type

$$x(1 - x)F + [C - (A + B + 1)x]F' - AB F = 0,$$

with parameters given by (to obtain the single-valued wave functions we must take a and b positive)

$$\begin{aligned} Z &= x^a(1 - x)^b {}_2F_1(A, B; C; x), & a &= \frac{-1 + \sqrt{1 + 4\lambda}}{4}, & b &= \frac{-1 + \sqrt{1 + 4\lambda}}{4}; \\ C &= 1 + \sqrt{\lambda + 1/4}, & A &= \frac{1}{2} + \sqrt{\lambda + \frac{1}{4}} + \sqrt{2\epsilon + 1}, & B &= \frac{1}{2} + \sqrt{\lambda + \frac{1}{4}} - \sqrt{2\epsilon + 1}. \end{aligned} \quad (71)$$

To obtain polynomials we must impose the following conditions:

$$\begin{aligned} \lambda + \frac{1}{4} &> 0, & 2\epsilon + 1 &> 0, & B &= \frac{1}{2} \\ &+ \sqrt{\lambda + \frac{1}{4}} - \sqrt{2\epsilon + 1} &= -N, & N &= 0, 1, 2, \dots \end{aligned}$$

from whence it follows the quantization rule

$$2\epsilon + 1 = \left(\frac{1}{2} + N + \sqrt{\lambda + \frac{1}{4}} \right)^2. \quad (72)$$

In particular, this formula gives

$$(2\epsilon + 1) - \left(\lambda + \frac{1}{4} \right) = \left(\frac{1}{2} + N \right)^2 + 2 \left(\frac{1}{2} + N \right) \sqrt{\lambda + \frac{1}{4}}, \quad (73)$$

with the notion

$$\left(\frac{1}{2} + N \right)^2 + 2 \left(\frac{1}{2} + N \right) \sqrt{\lambda + \frac{1}{4}} = \nu^2 > 0$$

it reads

$$(2\epsilon + 1) = \nu^2 + \left(\lambda + \frac{1}{4} \right) \quad (74)$$

which generalizes the known relation in flat space model: $2\epsilon = k^2 + \lambda$. The identity (74) permits us to express all parameters through ϵ, ν^2 :

$$\begin{aligned} C &= 1 + \sqrt{(2\epsilon + 1) - \nu^2}, \\ A &= \frac{1}{2} + \sqrt{(2\epsilon + 1) - \nu^2} + \sqrt{2\epsilon + 1}, \\ B &= \frac{1}{2} + \sqrt{(2\epsilon + 1) - \nu^2} - \sqrt{2\epsilon + 1}. \end{aligned} \quad (75)$$

However, it should be noted that such a form does not provide us with any advantage because quantization for the quantity $(2\epsilon + 1) - \nu^2$ is given by the differential equation for radial function $R(r)$. It is easily to obtain an operator with eigenvalues $\nu^2 + 3/4$:

$$\begin{aligned} \hat{\nu}^2 - \frac{3}{4} &= 2\hat{H}(r, \phi, z) - \hat{\lambda}(r), \\ \hat{\lambda}(r, \phi) &= -\frac{1}{\sin r} \frac{\partial}{\partial r} \sin r \frac{\partial}{\partial r} + \frac{1}{\sin^2 r} \left[-i \frac{\partial}{\partial \phi} + B(1 - \cos r) \right]^2, \end{aligned} \quad (76)$$

so that the operator is given by

$$\begin{aligned} \hat{\nu}^2(r, \phi, z) - \frac{3}{4} &= \left(\frac{1}{\cos^2 z} - 1 \right) \hat{\lambda}(r, \phi) - \frac{1}{\cos^2 z} \\ &\quad \times \frac{\partial}{\partial z} \cos^2 z \frac{\partial}{\partial z}. \end{aligned} \quad (77)$$

It commutes with Hamiltonian.

7 Presence of an electric field, the case of space S_3

Let us introduce into consideration an additional external electric field along the z axis

$$A_0 = E_0 \frac{\sin z}{\cos^2 z} \quad (78)$$

in the limit of flat space it becomes the uniform constant field E along z . As we will see below, this additional field does not destroy hypergeometric structure of the solution $Z(z)$. To take into account the presence of this field, it is sufficient to make one formal change in the previous calculations (in dimensionless notation)

$$\epsilon \implies \epsilon + eE_0 \frac{M\rho^2}{\hbar^2} \rho \frac{\sin z}{\cos^2 z} = \epsilon + \mu \frac{\sin z}{\cos^2 z}. \quad (79)$$

Also, one should use the identity

$$(2\epsilon + 2\mu \frac{\sin z}{\cos^2 z}) = 2\epsilon + 2\mu \frac{y}{1-y^2} = 2\epsilon + 2\mu \frac{2x-1}{4x(1-x)} = 2\epsilon + \frac{2\mu}{4} \left(-\frac{1}{x} + \frac{1}{1-x}\right).$$

Correspondingly, instead of eq. (68), we have

$$\begin{aligned} & x(1-x)F'' + [2a(1-x) - 2bx - 3x + 3]F' \\ & + \left[a(a-1) + \frac{3a}{2} - \frac{\lambda}{4} - \frac{2\mu}{4} \right] \frac{1}{x}F + \left[b(b-1) + 3b - \frac{3b}{2} - \frac{\lambda}{4} + \frac{2\mu}{4} \right] \frac{1}{1-x}F + \\ & + [-a(a-1) - ab - ab - b(b-1) - 3a - 3b + 2\epsilon]F = 0. \end{aligned} \quad (80)$$

By imposing two conditions

$$a^2 + \frac{a}{2} - \frac{\lambda + 2\mu}{4} = 0, \quad b^2 + \frac{b}{2} - \frac{\lambda - 2\mu}{4} = 0 \quad (81)$$

eq. (80) reduces to

$$x(1-x)F'' + [(2a + 3/2) - x(2a + 2b + 3)]F' - [(a + b + 1)^2 - 2\epsilon - 1]F = 0 \quad (82)$$

which is of hypergeometric type. Thus the solutions are given by

$$\begin{aligned} Z &= x^a(1-x)^b {}_2F_1(A, B; C; x), \quad x = \frac{1 + \sin z}{2}, \quad (1-x) = \frac{1 - \sin z}{2}, \\ a &= \frac{-1 + \sqrt{1 + 4(\lambda + 2\mu)}}{4}, \quad b = \frac{-1 + \sqrt{1 + 4(\lambda - 2\mu)}}{4}, \\ A &= \frac{1}{2} + \frac{1}{2}\sqrt{(\lambda + 2\mu) + \frac{1}{4}} + \frac{1}{2}\sqrt{(\lambda - 2\mu) + \frac{1}{4}} + \sqrt{2\epsilon + 1}, \\ B &= \frac{1}{2} + \frac{1}{2}\sqrt{(\lambda + 2\mu) + \frac{1}{4}} + \frac{1}{2}\sqrt{(\lambda - 2\mu) + \frac{1}{4}} - \sqrt{2\epsilon + 1}, \\ C &= 1 + \sqrt{(\lambda - 2\mu) + 1/4}. \end{aligned} \quad (83)$$

To obtain polynomials, we must impose the following restriction

$$(\lambda \pm 2\mu) + \frac{1}{4} > 0, \quad 2\epsilon + 1 > 0,$$

$$B = \frac{1}{2} + \sqrt{(\lambda + 2\mu) + \frac{1}{4}} + \sqrt{(\lambda - 2\mu) + \frac{1}{4}} - \sqrt{2\epsilon + 1} = -N, \quad N = 0, 1, 2, \dots$$

from whence it follows the quantization rule

$$2\epsilon + 1 = \left(\frac{1}{2} + N + \frac{1}{2}\sqrt{(\lambda + 2\mu) + \frac{1}{4}} + \frac{1}{2}\sqrt{(\lambda - 2\mu) + \frac{1}{4}} \right)^2 \quad (84)$$

where the quantization of λ is given by (61)). Thus, the energy levels are determined by two quantum numbers, (n, N) , and depends upon three parameters, curvature radius ρ , and magnitudes of fields: B and E .

References

- [1] E. Schrödinger. A method of determining quantum mechanical eigenvalues and eigenfunctions. Proc. R. Irish. Acad. A. **46**, 9 – 16 (1940).
- [2] L. Infeld. On a new treatment of some eigenvalue problems. Phys. Rev. **59**, 737 – 747 (1941).
- [3] L. Infeld and A. Schild. A note on the Kepler problem in a space of constant negative curvature. Phys. Rev. **67**, 121 – 122 (1945).
- [4] A.F. Stevenson. A note on the "Kepler problem" in a spherical space, and the factorization method of solving eigenvalue problems. Phys. Rev. **59**, 842 – 843 (1941).
- [5] P. Higgs, P. Dynamical symmetries in a spherical geometry. I. J. Phys. A: Math. Gen. **12**, 309 – 323 (1979).
- [6] H. Leemon. Dynamical symmetries in a spherical geometry. II. J. Phys. A: Math. Gen. **12**, 489 – 501 (1979).
- [7] Yu.A. Kurochkin, V.S. Otchik. Analogue of the Runge-Lenz vector and energy spectrum in Kepler problem on threedimensional sphere. Doklady AN BSSR. **23**, 987 – 990 (1979). (in Russian)
- [8] A.A. Bogush, Yu.A. Kurochkin, V.S. Otchik. On quantummechanical Kepler problem in Lobachevsky space. Doklady AN BSSR. **24**, 19 – 22 (1980). (in Russian)
- [9] V.S. Otchik, V.M. Red'kov. *Quantum-mechanical Kepler problem in spaces of constant curvature*. Preprint no. 298. (Institute of Physics, Minsk, 1983). 50 Pp. (in Russian)
- [10] A.A. Izmetiev. Exactly solvable potential model for quarkoniums. Yadernay Fizika. **52**, 1697 – 1710 (1990). (in Russian)
- [11] V.I. Murzov, V.V. Kudriashov, L.M. Tomil'chik. *Triplet state for 2-fermion system with uniform relative space*. Preprint no. 497. (Institute of Physics, Minsk, 1988). 34 Pp. (in Russian)
- [12] V.I. Murzov, V.V. Kudriashov, L.M. Tomil'chik. *Quantum-mechanical composite systems with non-euclidean relative space*. Preprint no. 568. (Institute of Physics, Minsk, 1989). 20 Pp. (in Russian)
- [13] Yu.A. Kurochkin. Non-Euclidean quantum mechanics for exciton in semiconducting quantum dots. Doklady AN BSSR. **38**, 36 – 40. (1994)(in Russian).
- [14] V. Gritzev, Yu. Kurochkin. Model of excitations in quantum dots based on quantum mechanics in spaces of constant curvature. Phys. Rev. B. **64**, no. 3, 035308 (2001).
- [15] M.N. Olevsky. Three-orthogonal coordinate systems in spaces of constant curvature, in which equation $\Delta_2 U + \lambda U = 0$ permits the full separation of variables. Mathematical collection. **27**, 379 – 426 (1950). (in Russian).
- [16] L. Landau. Diamagnetismus der Metalle. Zeit. Phys. **64**, 629 – 637 (1930).
- [17] L.D. Landau, E.M. Lifshitz. *The theory of field*. (Moscow, 1973). (in Russian)

- [18] L.D. Landau, E.M. Lifshitz. *Quantum mechanics, nonrelativistic theory*. (Moscow, 1974). (in Russian)
- [19] J. R. Klauder, E. Onofri. Landau Levels and Geometric Quantization. *Int. J.Mod. Phys. A.* P. 3939-3949 (1989).
- [20] Klauder, E. Onofri J.E. Avron, A. Pnueli. Landau Hamiltonians on Symmetric Spaces. In: *Ideas and methods in mathematical analysis, stochastics, and applications*. Vol. II. S. Alverio et al., eds. (Cambridge Univ. Press, Cambridge, 1992).
- [21] G. V. Dunne. Hilbert Space for Charged Particles in Perpendicular Magnetic Fields. *Ann. Phys. (N.Y.)* **215**, 233 – 263 (1992).
- [22] J. Negro, M.A. del Olmo, A. Rodríguez-Marco. Landau quantum systems: an approach based on symmetry. *arXiv:quantum-ph/0110152*.
- [23] Nadav Drukker, Bartomeu Fiol, Joan Simón. Gödel-type Universes and the Landau problem. *hep-th/0309199*.
- [24] J. Gamboa, M. Loewe, F. Mendez, J. C. Rojas Noncommutative Quantum Mechanics: The Two-Dimensional Central Field. *Int. J. Mod. Phys. A.* **17**, 2555 – 2566 (2002).
- [25] J. Gamboa, M. Loewe, F. Mendez, J. C. Rojas The Landau problem and noncommutative quantum mechanics. *Mod. Phys. Lett. A.* **16**, 2075 – 2078 (2001).
- [26] J. Gamboa, M. Loewe, J.C. Rojas. Non-Commutative Quantum Mechanics. *Phys. Rev. D.* **64**, 067901 (2001).
- [27] H. Bateman, A. Erdélyi. *Higher transcendental functions*. Vol. I. (New York, McGraw-Hill) 1953.